

# Sandblasting The R-process: Spallation Of R-process Nuclei Ejected From A NSNS Event

Xilu Wang,<sup>1</sup> Brian Fields,<sup>2</sup> Matthew Mumpower,<sup>3</sup>

Trevor Sprouse,<sup>1</sup> Rebecca Surman,<sup>1</sup> and Nicole Vassh<sup>1</sup>

<sup>1</sup>*University of Notre Dame, Notre Dame, IN 46556, United States*

<sup>2</sup>*University of Illinois at Urbana-Champaign, Urbana, IL 61801, United States*

<sup>3</sup>*Los Alamos National Lab, Los Alamos, NM 87544, United States*

Neutron star mergers are r-process nucleosynthesis sites, which eject materials at high velocity ranging from 0.1c to 0.3c for different regions. Thus the r-process nuclei ejected from a neutron star merger event are sufficiently energetic to have spallation nuclear reactions with the interstellar medium particles. The spallation reactions tend to shift the r-process abundance patterns towards the solar data, and smooth the abundance shapes. The spallation effects depend on both the initial r-process nuclei conditions, which is determined by the astrophysical trajectories and nuclear data adopted for the r-process nucleosynthesis, and the propagation with various ejecta velocities and spallation cross-sections.