

# On Barium stars and the s-process in AGB stars

Borbála Cseh,<sup>1</sup> Maria Lugaro,<sup>1</sup> Valentina D’Orazi,<sup>2</sup> Denise B. de Castro,<sup>3</sup> Claudio B. Pereira,<sup>4</sup> Amanda I. Karakas,<sup>5</sup> László Molnár,<sup>1</sup> Emese Plachy,<sup>1</sup> and Róbert Szabó<sup>1</sup>

<sup>1</sup>*Konkoly Observatory, Research Centre for Astronomy and Earth Sciences,  
Hungarian Academy of Sciences, H-1121 Budapest, Hungary*

<sup>2</sup>*INAF Osservatorio Astronomico di Padova,  
vicolo dell’Osservatorio 5, 35122 Padova, Italy*

<sup>3</sup>*Nicolaus Copernicus Astronomical Centre of the  
Polish Academy of Sciences, 00-716 Warsaw, Poland*

<sup>4</sup>*Observatorio Nacional, Rua General Jos Cristino,  
77 Sao Cristovao, Rio de Janeiro, Brazil*

<sup>5</sup>*Monash Centre for Astrophysics, School of Physics and Astronomy,  
Monash University, VIC 3800, Australia*

Barium stars belong to a binary system where the companion star has evolved through the AGB phase and transferred elements heavier than Fe produced by the slow neutron-capture process onto the secondary star, which is now observed. A new large set of homogeneous high resolution spectra of Ba stars makes it now possible to meaningfully compare the observational data with different AGB models and with other observations (e.g., post-AGB stars or stardust grains). The Ba star data shows an incontestable increase of the hs-type/ls-type element ratio (for example, [Ce/Y]) with decreasing the metallicity. The trend in the Ba star observations is predicted by non-rotating AGB models where  $^{13}\text{C}$  is the main neutron source. Observations of the cores of red giant stars and of white dwarfs (the ancestors and the progeny of AGB stars, respectively) inferred via asteroseismology from Kepler observations show low core rotational velocities, which is in agreement with the results from the Ba star data and may derive from coupling between the core and the envelope.